

Goat anti-Aconitase 2 (aa541-555), Biotinylated Antibody

Item Number	dAP-3439
Target Molecule	Principle Name: Aconitase 2 (aa541-555), Biotinylated; Official Symbol: ACO2; All Names and Symbols: ACO2; aconitase 2; ACONM; HEL-S-284; ICRD; OCA8; OPA9; aconitase 2, mitochondrial; citrate hydrolyase; epididymis secretory sperm binding protein Li 284; mitochondrial aconitase; Accession Number (s): NP_001089.1; Human Gene ID(s): 50; Non-Human GeneID(s):
Immunogen	CQDTYQHPPKDSSGQH., is from internal region
Applications	Pep ELISA, WB, Species Tested: Human, Mouse, Rat, Pig
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 16000.
Western Blot	Western Blot: Approx 90kDa band observed in Human Heart lysates, in Human, Mouse and Rat Adipose lysates and in Mouse, Rat and Pig Skeletal Muscle lysates (calculated MW of 85.4kDa according to NP_001089.1). Recommended concentration: 0.01-0.03µg/ml. See
IHC	
Reference	Reference(s): Bota DA, Davies KJ. Lon protease preferentially degrades oxidized mitochondrial aconitase by an ATP-stimulated mechanism. Nature cell biology 2002 Sep 4 (9): 674-80..PMID: 12198491->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**