



## soluble CD40 Ligand/TRAP Human Recombinant, His Tag

Item Number rAP-0111

CD40-L, Tumor necrosis factor ligand superfamily member 5, TNF-related activation protein, TRAP, T cell Synonyms

antigen Gp39, CD154 antigen, sCD40, IGM, IMD3, HIGM1, T-BAM, TNFSF5, hCD40L, CD40LG.

Description CD40LG Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a

single, non-glycosylated, polypeptide chain containing 169 amino acids (113-261 a.a.) and having a molec-

ular mass of 18.3kDa. The CD40LG is purified by proprietary chromatographic techniques.

P29965 **Uniprot Accesion Number** 

MGSSHHHHHH SSGLVPRGSH MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYYTMSNNL **Amino Acid Sequence** 

VTLENGKQLT VKRQGLYYIY AQVTFCSNRE ASSQAPFIAS LCLKSPGRFE RILLRAANTH

SSAKPCGQQS IHLGGVFELQ PGASVFVNVT DPSQVSHGTG FTSFGLLKL.

Escherichia Coli. Source

**Physical Appearance** 

and Stability

Sterile Filtered clear solution. Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Formulation and Purity The CD40LG solution (0.5mg/ml) 20mM Tris-HCl buffer (pH 8.0), 1mM DTT, 40% glycerol and 0.2M NaCl.

Greater than 95.0% as determined by SDS-PAGE.

**Application** 

Solubility

**Biological Activity** 

Shipping Format and Condition Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only