



Fibroblast Growth Factor Receptor 4 Fc Chimera Human Recombinant

Item Number rAP-1108

Synonyms Fibroblast Growth Factor Receptor 4, EC 2.7.10.1, JTK2, TKF, Tyrosine Kinase Related To Fibroblast

Growth Factor Receptor, Hydroxyaryl-Protein Kinase, Protein-Tyrosine Kinase, Tyrosylprotein Kinase,

CD334 Antigen, EC 2.7.10, FGFR-4, CD334, FGFR4.

Description Soluble FGFR-4a (IIIc) Fc Chimera Human Recombinant fused with Xa cleavage site with the Fc part of

human IgG1 produced in baculovirus is a heterodimeric, glycosylated, Polypeptide chain and having a mo-

lecular mass of 170 kDa. The FGFR4 is purified by proprietary chromatographic techniques.

Uniprot Accesion Number P22455

Amino Acid Sequence

Source Insect Cells.

Physical Appearance

and Stability

Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized FGFR4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FGFR4 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to

add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Formulation and Purity CD334 was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. Greater than

90.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Application

Solubility It is recommended to reconstitute the lyophilized FGFR-4 in sterile PBS not less than 100 μg/ml, which can

then be further diluted to other aqueous solutions.

Biological Activity Determined by its ability to inhibit human FGF acidic-dependent proliferation on R1 cells. The ED50 for this

effect is typically at 15.0-30.0 ng/ml.

Shipping Format and Condition Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only