



Vascular Endothelial Growth Factor Receptor-1 D4 Human Recombi-

Item Number rAP-2112

FLT-1, FLT1, Tyrosine-protein kinase receptor FLT, Flt-1, Tyrosine-protein kinase FRT, Fms-like tyrosine Synonyms

kinase 1, VEGFR-1.

Description Soluble FLT1 D1-4 Human Recombinant produced in baculovirus is monomeric, glycosylated, polypeptide

containing 457 amino acids and having a molecular mass of 55 kDa. The soluble receptor protein contains only the first 4 extracellular domains, which contain all the information necessary for binding of VEGF. The

P17948 **Uniprot Accesion Number**

Amino Acid Sequence

Source Insect Cells.

Physical Appearance

and Stability

Sterile Filtered White lyophilized (freeze-dried) powder. Lyophilized FLT-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FLT1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a

carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

FLT1 D1-4 was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. Greater Formulation and Purity

than 90.0% as determined by(a)Analysis by RP-HPLC.(b)Analysis by SDS-PAGE.

Application

Solubility It is recommended to reconstitute the lyophilized FLT1 D4 in sterile water not less than 100 µg/ml, which

can then be further diluted to other aqueous solutions.

The activity of FLT1D1-4 was determined by its ability to abolish the binding of iodinated VEGF to solid **Biological Activity**

surfaces or cell surfaces, and in Far-Western and cross-linking experiments with iodinated VEGF.

Shipping Format and Condition Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only