

Intercellular Adhesion Molecule-2 Human Recombinant

Item Number	rAP-3496
Synonyms	Intercellular Adhesion Molecule 2, ICAM2, ICAM-2, CD102, CD102 Antigen.
Description	ICAM2 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 225 amino acids (22-223) and having a molecular mass of 24.8 kDa. ICAM2 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.
Uniprot Accession Number	P13598
Amino Acid Sequence	MGSSHHHHHH SSGLVPRGSH MGSSDEKVFE VHVPRPKKLAV EPKGSLEVNC STTCNQPEVG GLETSLDKIL LDEQAQWKHY LVSNIHDTV LQCHFTCSGK QESMNSNVSV YQPPRQVILT LQPT- LVAVGK SFTIECRVPT VEPLDSLTLF LFRGNETLHY ETFGKAAPAP QEATATFNST ADREDGHRNF SCLAVLDLMS RGGNIFHKHS APKMLEIYEP VSDSQ.
Source	Escherichia Coli.
Physical Appearance and Stability	Sterile Filtered colorless solution. Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
Formulation and Purity	The ICAM2 solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 10% glycerol and 1mM DTT. Greater than 90.0% as determined by SDS-PAGE.
Application	
Solubility	
Biological Activity	
Shipping Format and Condition	Lyophilized powder at room temperature.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**