

Mouse Monoclonal Antibody to SMC1

Catalogue Number	sAP-0346
Target Molecule	Name: SMC1 Aliases: SMC1; SMCB; CDLS2; SB1.8; SMC1L1; DXS423E; KIAA0178; MGC138332; SMC1alpha; DKFZp686L19178; SMC1A MW: 143kDa
Description	Proper cohesion of sister chromatids is a prerequisite for the correct segregation of chromosomes during cell division. The cohesin multiprotein complex is required for sister chromatid cohesion. This complex is composed partly of two structural maintenance of chromosomes (SMC) proteins, SMC3 and either SMC1L2 or the protein encoded by this gene. Most of the cohesin complexes dissociate from the chromosomes before mitosis, although those complexes at the kinetochore remain. Therefore, the encoded protein is thought to be an important part of functional kinetochores. In addition, this protein interacts with BRCA1 and is phosphorylated by ATM, indicating a potential role for this protein in DNA repair. This gene, which belongs to the SMC gene family, is located in an area of the X-chromosome that escapes X inactivation.
Immunogen	Purified recombinant fragment of human SMC1 expressed in E. Coli. ;
Reactive Species	Human;
Clone	MM5B6;
Size and Concentration	100µg/1mg/ml
Supplied as	Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide
Reconstitution/Storages	Reconstituted with 100µl sterile DI H ₂ O, at stored at 4°C or -20°C for short or long term storage
Applications	ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000; ICC: 1 to 200 - 1 to 1000; FCM: 1 to 200 - 1 to 400
Shipping	Regular FEDEX overnight shipment (ambient temperature)
Reference	1. Cell Cycle. 2006 Nov 1;5(21):2537-42. ; 2. FEBS Lett. 2007 Jun 26;581(16):3005-12.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**