



Mouse Monoclonal Antibody to CD59

Cataloge Number sAP-0775

Target Molecule Name: CD59

Aliases: 1F5; EJ16; EJ30; EL32; G344; MIN1; MIN2; MIN3; MIRL; HRF20; MACIF; MEM43; MIC11;

MSK21; 16.3A5; HRF-20; MAC-IP; p18-20

MW: 14.2kDa

Descrption This gene encodes a cell surface glycoprotein that regulates complement-mediated cell lysis, and it is in-

volved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. This protein also plays a role in signal transduction pathways in the activation of T cells. Mutations in this gene cause CD59 deficiency, a disease resulting in hemolytic anemia and thrombosis, and which causes cerebral infarction. Multiple alternatively spliced transcript variants, which encode the same

protein, have been identified for this gene.;;

Immunogen Purified recombinant fragment of human CD59 (AA: 31-111) expressed in E. Coli.

Recitative Species Human;

Clone MM8D2B8;

Size and Concentration 100µg/1mg/ml

Supplied as Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide

Reconstitution/Storages Reconstitued with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage

Applications ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000; ICC: 1 to 50 -1 to 200; FCM: 1 to

200 - 1 to 400

Shipping Regular FEDEX overnight shipment (ambient temperature)

Reference 1.Cell Immunol. 2010;265(2):127-32.; 2.Chin Med J (Engl). 2009 Sep 20;122(18):2123-8.;

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only