

# Anti-Mouse CD31 PE-Cyanine7

Catalog Number: 03412-77

RUO: For Research Use Only. Not for use in diagnostic procedures.

### **Product Information**

**Clone:** 390

Format/Conjugate: PE-Cyanine7 Concentration: 0.2 mg/mL

**Reactivity:** Mouse **Laser:** Blue (488nm)

**Peak Excitation:** Not Applicable **Peak Excitation:** Not Applicable

Filter: Not Applicable

Brightness (1=dim,5=brightest): Not Applicable

Isotype: Rat IgG2a, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

**Applications:** FC

### **Description**

The 390 monoclonal antibody specifically reacts with mouse cd31, a 130-140 kDA type I transmembrane glycoprotein also known as platelet-endothelial cell adhesion molecule-1 (PECAM-1). CD31 is reported to bind to CD38 and is expressed on platelets, monocytes, granulocytes, and endothelial cells. It plays a role in angiogenesis, wound healing, cellular migration, and signal transduction.

## **Preparation & Storage**

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

## **Application Notes**

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.5$  ug per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

#### References

1.HORENSTEIN, A., STOCKINGER, H., Imhof, B., MALAVASI, F. (1998). CD38 binding to human myeloid cells is mediated by mouse and human CD31.Biochem. J,;330, 1129-1135.

- 2. Baldwin, H. S., Shen, H. M., Yan, H. C., DeLisser, H. M., Chung, A., Mickanin, C., ... Albelda, S. M. (1994). Platelet endothelial cell adhesion molecule-1 (PECAM-1/CD31): alternatively spliced, functionally distinct isoforms expressed during mammalian cardiovascular development.;Development.;120(9), 2539-2553.
- 3. DeLisser, H. M., Christofidou-Solomidou, M., Strieter, R. M., Burdick, M. D., Robinson, C. S., Wexler, R. S., ... Albelda, S. M. (1997). Involvement of endothelial PECAM-1/CD31 in angiogenesis.;The American journal of pathology,;151(3), 671.