



# Anti-Human CD24 FITC

Catalog Number: 06321-50

RUO: For Research Use Only. Not for use in diagnostic procedures.

### **Product Information**

Clone: ML5

Format/Conjugate: FITC

Concentration: 5uL(2.0ug)/test

Reactivity: Human
Laser: Blue (488nm)
Peak Emission: 520nm
Peak Excitation: 494nm

Filter: 530/30

Brightness (1=dim,5=brightest): 3

Isotype: Mouse IgG2a, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.

**Storage:** Product should be kept at 2-8°C.

**Applications:** FC

### **Description**

The ML5 monoclonal antibody specifically reacts with human CD24, a 35-45 kDA molecule also known as the Heat Stable Antigen (HAS), Ly-52, Nectadrin. It can be used as a marker for distinguishing between lymphocyte developmental stages as its expression varies on T and B cells during differentiation. CD24 is also expressed on monocytes, dendritic cells, hematopoietic stem cells, epidermal Lagershans cells, and neurons.

## **Preparation & Storage**

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

## **Application Notes**

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. The antibody can be used at less than or equal to 5  $\mu$ L per test. A test is the amount of antibody required to stain a cell sample in the final volume of 100  $\mu$ L.

#### References

- 1.McMichael, A. J. (1987).;Leucocyte typing III: white cell differentiation antigens. Oxford University Press, USA.
- 2. Sagiv, E., Kazanov, D., Arber, N. (2006). CD24 plays an important role in the carcinogenesis process of the pancreas.; Biomedicine pharmacotherapy, 60(6), 280-284.
- 3. Kristiansen, G., Winzer, K. J., Mayordomo, E., Bellach, J., Schlüns, K., Denkert, C., ... Dietel, M. (2003). CD24 expression is a new prognostic marker in breast cancer.; Clinical Cancer Research,; 9(13), 4906-4913.

4.;