

Anti-Human CD45 Biotin

Catalog Number :07111-30

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Clone: HI30

Format/Conjugate: Biotin

Concentration: 0.5 mg/mL

Reactivity: Human

Laser: Not Applicable

Peak Emission: Not Applicable

Peak Excitation: Not Applicable

Filter: Not Applicable

Brightness (1=dim,5=brightest): Not Applicable

Isotype: Mouse IgG1, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, pH7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

Applications: FC

Description

The HI130 monoclonal antibody specifically reacts with the 180 kDa, 190 kDa, 205 kDa, and 220 kDa isoforms of the human leukocyte common antigen (LCA) CD45. It is expressed on lymphocytes, granulocytes, monocytes, thymocytes, and eosinophils, but not on mature erythrocytes, platelets, mature erythroid cells of bone marrow, and non-hematopoietic tissues. CD45 is essential for T cell activation and the tyrosine phosphatase activity of its intracellular region is integral for signal transduction.

Preparation & Storage

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

Application Notes

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. For flow cytometric staining, the suggested use of this reagent is ≤0.25 ug per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

References

1. Knapp W;(1989) Leucocyte typing IV: white cell differentiation antigens. Oxford University Press, 1989.
2. Ninomiya, M., Abe, A., Katsumi, A., Xu, J., Ito, M., Arai, F., ... Naoe, T. (2006). Homing, proliferation and survival sites of human leukemia cells in vivo in immunodeficient mice.;*Leukemia*,;21(1), 136-142.
3. Yoshino, N., AMI, Y., TERAOKA, K., TASHIRO, F., HONDA, M. (2000). Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies.;*Experimental Animals*,;49(2), 97-110.