

# Anti-Human CD8a PE

Catalog Number:10111-60

RUO: For Research Use Only. Not for use in diagnostic procedures.

#### **Product Information**

Clone: Hit8a

Format/Conjugate: PE

Concentration: 5 uL (0.06 ug)/test

Reactivity: Human Laser: Blue (488nm) Peak Emission: 578nm Peak Excitation: 496nm

Filter: 585/40

**Brightness (1=dim,5=brightest):** 5

Isotype: Mouse IgG1, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

**Applications:** FC

#### **Description**

The Hit8a monoclonal antibody reacts with the human CD8a molecule, a 32 kDa cell surface receptor expressed either as a heterodimer (CD8  $\alpha/\beta$ ) or as a homodimer (CD8  $\alpha/\alpha$ ) on the majority of thymocytes, a subpopulation of mature T cells, and natural killer cells. CD8 interacts with the major histocompatibility complex class I (MHC class I) molecules on antigenpresenting cells or epithelial cells. The Hit8a antibody reacts with 13-48% of peripheral lymphocytes, 80% of thymocytes, and a subset of natural killer cells.

HIT8a, RPA-T8, and OKT8 antibodies do not compete with each other for binding to peripheral leukocytes, meaning that that they do not recognize the same epitope or block each other by steric hindrance.

## **Preparation & Storage**

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

## **Application Notes**

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. The antibody can be used at less than or equal to 5  $\mu$ L per test. A test is the amount of antibody required to stain a cell sample in the final volume of 100  $\mu$ L.

### References

- 1. Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York
- 2. Barclay, A. N., Brown, M. H., Law, S. A. K. A., McKnight, A. J., Tomlinson, M. G., van der Merwe, P. A. (1997).; The leucocyte antigen factsbook. Academic Press.