

Anti-Caspase-8/CASP8 Antibody Picoband™

Catalog Number: A00042-3

About CASP8

CASP8 is also known as CAP4, MACH or MCH5. This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. In addition, this protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined.

Overview

Product Name	Anti-Caspase-8/CASP8 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Caspase-8/CASP8 Antibody Picoband™ catalog # A00042-3. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q14790

Technical Details

Immunogen	E.coli-derived human Caspase-8/CASP8 recombinant protein (Position: S217-D479).
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5 ug/ml, Human Flow Cytometry, 1-3 ug/1x10⁶ cells, Human Direct ELISA, 0.1-0.5 ug/ml, Human

Anti-Caspase-8/CASP8 Antibody Picoband™ (A00042-3) Images




Figure 1. Western blot analysis of Caspase-8/CASP8 using anti-Caspase-8/CASP8 antibody (A00042-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human Jurkat whole cell lysates,

Lane 3: human Raji whole cell lysates,

Lane 4: human K562 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Caspase-8/CASP8 antigen affinity purified polyclonal antibody (Catalog # A00042-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Caspase-8/CASP8 at approximately 55 kDa. The expected band size for Caspase-8/CASP8 is at 55 kDa.




Figure 2. Flow Cytometry analysis of K562 cells using anti-Caspase-8/CASP8 antibody (A00042-3).

Overlay histogram showing K562 cells stained with A00042-3 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Caspase-8/CASP8 Antibody (A00042-3, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

3 Publications Citing This Product

1. PubMed ID: 10.1248/bpb.b13-00794, Involvement of substance P/neurokinin-1 receptor in the analgesic and anticancer activities of minimally toxic fraction from the traditional Chinese medicine Liu-Shen-Wan in vitro
2. PubMed ID: 10.3892/or.2015.4530, Redox cycling of a copper complex with benzaldehyde nitrogen mustard-2-pyridine carboxylic acid hydrazone contributes to its enhanced antitumor activity, but no change in the mechanism of action occurs after chelation
3. PubMed ID: 10.3892/ijo.2015.3039, Autophagy is involved in recombinant Newcastle disease virus (rL-RVG)-induced cell death of stomach adenocarcinoma cells in vitro

Visit bosterbio.com/anti-caspase-8-casp8-picoband-trade-antibody-a00042-3-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Caspase-8/CASP8 Antibody