

Anti-Smad2 (Phospho-S255) Antibody

Catalog Number: A00090S255

About SMAD2

Beta amyloid, often abbreviated as A-beta, is a protein that builds up in the brains of persons with Alzheimer's disease, collecting in clumps called plaques or senile plaques. While some researchers question whether beta amyloid is the cause of the dementia, most agree that it is involved in the disruption of thinking that is a hallmark of the disease. In some cases of familial Alzheimer's disease, mutations in genes for the proteins called the presenilins lead to increased production of amyloid. Researchers have been looking at how presenilin-1 in particular contributes to the excess buildup of beta amyloid. Presenilin-1 apparently acts to increase the activity of gamma-secretase, an enzyme that changes a normal protein (amyloid precursor protein or APP) into beta amyloid itself. Furthermore, presenilin-1 might be gamma-secretase. Anti-Beta Amyloid Antibody is useful for researchers interested in TLR signaling and Alzheimer's research.

Overview

Product Name	Anti-Smad2 (Phospho-S255) Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Smad2 (Phospho-S255) Antibody catalog # A00090S255. Tested in WB,IHC,IP applications. This antibody reacts with Human,Mouse,Rat.
Application	IP, IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q15796

Technical Details

Immunogen	Synthesized peptide derived from human Rad51 around the phosphorylation site of Y315.
Predicted Reactive Species	Canine, Monkey
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	ProA affinity purified
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:1,000 IHC: 1:50-1:200

Anti-Smad2 (Phospho-S255) Antibody (A00090S255) Images




Figure 1. Immunohistochemistry validation of SMAD2 using Anti-Smad2 (Phospho-S255) Antibody (A00090S255).

Immunohistochemical analysis of paraffin-embedded mouse uterus tissue using anti-Phospho-Smad2(S255) antibody. Counter stained with hematoxylin. For more protocol information of IHC

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Smad2 (Phospho-S255) Antibody