

Anti-SMAD2 Antibody

Catalog Number: A00090

About SMAD2

Smad2 (also known as Mothers against decapentaplegic homolog 2, Mothers against DPP homolog 2, Mad2, hMAD-2 or hSMAD2) is a member of the Smad family of proteins which are similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively, spliced transcript variants have been observed for this gene. SMAD2 may act as a tumor suppressor in colorectal carcinoma. It positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator. SMAD2 may be associated with diseases such as Keloids and Ureteral Disease. Anti-SMAD2 Antibody is useful for researchers interested in SMAD pathways, transcription factor activity, sequence-specific DNA binding, and cancer research.

Overview

Product Name	Anti-SMAD2 Antibody
Reactive Species	Human, Monkey
Description	Boster Bio Anti-SMAD2 Antibody (Catalog # A00090). Tested in ELISA, WB applications. This antibody reacts with Human, Monkey.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	Q15796

Technical Details

Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated
-----------	---

	immunizations with a synthetic peptide corresponding to an internal region of human Smad2 protein.
Predicted Reactive Species	Bovine, Canine, Monkey
Isotype	IgG
Form	Liquid (sterile filtered)
Concentration	0.55 mg/mL by UV absorbance at 280 nm
Purification	This affinity purified antibody is directed against human Smad2 protein. The product was affinity purified from monospecific antiserum by immunoaffinity chromatography. A BLAST analysis was used to suggest cross-reactivity with Smad2 protein from human, mouse and rat based on 100% homology with the immunizing sequence. Reactivity against homologues from other sources is not known. Also, the antibody is Smad2 specific, and reactivity to other Smad proteins (specifically Smad1, Smad3, Smad4, and Smad7) is not detected in over-expressed cell lysates (Personal Communication, Kathleen Flanders, CCR-NCI, Bethesda, MD).
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>ELISA: 1:100,000</p> <p>IP: User optimized</p> <p>WB: 1:1,000 - 1:3,000</p>

Anti-SMAD2 Antibody (A00090) Images




Figure 1. Western blot analysis of SMAD2 using anti-SMAD2 antibody (A00090). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SMAD2 antigen affinity purified polyclonal antibody (Catalog # A00090) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-Rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # SA1022) with Tanon 5200 system. A specific band was detected for SMAD2.

3 Publications Citing This Product

1. PubMed ID: 27019660, Effect of Kuijie Granule on the Expression of TGF- β /Smads Signaling Pathway in Patients with Ulcerative Colitis
2. PubMed ID: 26261569, Astragaloside effect on TGF- β 21, SMAD2/3, and β 1-SMA expression in the kidney tissues of diabetic KKAY mice
3. PubMed ID: 30090338, Mouse hepatic neoplasm formation induced by trace level and low frequency exposure to diethylnitrosamine through β 2-catenin signaling pathway

Visit bosterbio.com/anti-smad2-antibody-a00090-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SMAD2 Antibody