

Anti-Argonaute 2 AGO2 Antibody

Catalog Number: A00189-1

About AGO2

FLT3 encodes a class III receptor tyrosine kinase that regulates hematopoiesis. The receptor consists of an extracellular domain composed of five immunoglobulin-like domains, one transmembrane region, and a cytoplasmic kinase domain split into two parts by a kinase-insert domain. The receptor is activated by binding of the fms-related tyrosine kinase 3 ligand to the extracellular domain, which induces homodimer formation in the plasma membrane leading to autophosphorylation of the receptor. The activated receptor kinase subsequently phosphorylates and activates multiple cytoplasmic effector molecules in pathways involved in apoptosis, proliferation, and differentiation of hematopoietic cells in bone marrow. Mutations that result in the constitutive activation of this receptor result in acute myeloid leukemia and acute lymphoblastic leukemia.

Sekine S et.al. (2008) J Immunol. ;180(12):8126-34

Pratz K et.al. (2008) Leuk Lymphoma. 2008 ;49(5):852-63.

Al Shaer L et.al. (2008) Br J Haematol. 141(4):483-93.

Overview


Product Name	Anti-Argonaute 2 AGO2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Argonaute 2 AGO2 Antibody catalog # A00189-1. Tested in WB,ICC/IF,IHC,IP,Flow Cytometry applications. This antibody reacts with Human,Mouse,Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UKV8

Technical Details


Immunogen	Synthesized peptide derived from human Flt-3 around the phosphorylation site of Y969.
Predicted Reactive Species	Canine, Monkey
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG

Form	Liquid
Concentration	1 mg/ml
Purification	ProA affinity purified
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:1,000-1:2,000 ICC: 1:50-1:200 IHC: 1:50-1:200 FC: 1:50-1:100

Anti-Argonaute 2 AGO2 Antibody (A00189-1) Images



ICC staining Argonaute 2 in HeLa cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Western blot analysis of Argonaute 2 on different lysates using anti-Argonaute 2 antibody at 1/1,000 dilution. Positive control: Lane 1: HeLa Lane 2: MCF-7

1 Publications Citing This Product

1. PubMed ID: 33177098, Xin X, Kumar V, Lin F, Kumar V, Bhattarai R, Bhatt VR, Tan C, Mahato RI. Redox-responsive nanoplatfrom for codelivery of miR-519c and gemcitabine for pancreatic cancer therapy. Sci Adv.2020 Nov 11;6(46):eabd6764.doi:10.1126/sciadv.abd6764.PMID:33177098.

Visit bosterbio.com/anti-argonaute-2-ago2-antibody-a00189-1-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Argonaute 2 AGO2 Antibody