

Anti-ATR Antibody

Catalog Number: A00262

About ATR

Ataxia Telangiectasia Mutated (ATM) and Rad 3-related protein (ATR) is a phosphatidyl-inositol kinase (PK)-related kinase which functions in response to DNA damage and repair as well as at DNA replication checkpoints during the cell cycle. ATR activates checkpoint signaling upon genotoxic stresses, such as ionizing radiation (IR), ultraviolet light (UV), or DNA replication stalling, thereby acting as a DNA damage sensor. ATR is a member of the DNA-PK kinase family and is closely related to ATM and DNA-PK for which DNA stimulates the observed kinase activity.

Chromosomal remodeling proteins have also been reported to associate with ATR complexes, including histone deacetylases (HDAC1, HDAC2 and CHD4). ATR is known to phosphorylate BRCA1, CHEK1, MCM2, RAD17, RPA2, SMC1 and TP53/p53 which collectively inhibit DNA replication and mitosis and promote DNA repair, recombination and apoptosis. ATR is a nuclear protein, but can also be found in PML nuclear bodies in certain cell types. ATR is recruited to chromatin during S-phase and redistributes to discrete nuclear foci upon DNA damage, hypoxia or replication fork stalling.

Overview

Product Name	Anti-ATR Antibody
Reactive Species	Human
Description	Boster Bio Anti-ATR Antibody (Catalog # A00262). Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	Q13535

Technical Details

Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human ATR protein.
Predicted Reactive Species	Canine, Fish, Monkey, Xenopus
Cross Reactivity	No cross reactivity with other proteins.








Isotype	IgG
Form	Liquid (sterile filtered)
Concentration	1.28 mg/mL by UV absorbance at 280 nm
Purification	This product was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody is specific for human ATR protein. A BLAST analysis was used to suggest cross-reactivity with ATR from human, mouse, rat, monkey, dog, fish and Xenopus sources based on a 100% homology with the immunizing sequence. Cross-reactivity with ATR from other sources has not been determined.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: ELISA: 1:15,000 - 1:70,000 WB: 1:1,000 - 1:5,000



Anti-ATR Antibody (A00262) Images



Western blot analysis of ATR expression in HELA cell nuclear extract (lane 1), MW markers (Lane M) and negligible staining after pre-incubation of antibody with the immunizing peptide (50 ug peptide for 1 h at room temperature followed by centrifugation) (lane 2). ATR at 301KD was detected using rabbit anti-ATR Antigen Affinity purified polyclonal antibody (Catalog # A00262) at 1:1400. The blot was developed using chemiluminescence (ECL) method (Catalog # EK1002).

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-ATR Antibody