

Anti-LPL Antibody (Center)

Catalog Number: A00316-1

About LPL

LPL encodes lipoprotein lipase, which is expressed in heart, muscle, and adipose tissue. LPL functions as a homodimer, and has the dual functions of triglyceride hydrolase and ligand/bridging factor for receptor-mediated lipoprotein uptake. Severe mutations that cause LPL deficiency result in type I hyperlipoproteinemia, while less extreme mutations in LPL are linked to many disorders of lipoprotein metabolism. [provided by RefSeq].

Overview

Product Name	Anti-LPL Antibody (Center)
Reactive Species	Human
Description	Boster Bio Anti-LPL Antibody (Center) (Catalog # A00316-1). Tested in WB, IHC-P, Flow Cytometry application(s). This antibody reacts with Human.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P06858

Technical Details

Immunogen	This LPL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 300-327 amino acids from the Central region of human LPL.
Predicted Reactive Species	Bovine, Mouse, Pig, Rat, Sheep
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

WB: 1:1000-1:2000


IHC-P: 1:10-1:50

FC: 1:25

Anti-LPL Antibody (Center) (A00316-1) Images



LPL Antibody (Center) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the LPL antibody detected the LPL protein (arrow).



All lanes : Anti-LPL Antibody (Center) at 1:1000-1:2000 dilution

Lane 1: MCF-7 whole cell lysate

Lane 2: HL-60 whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary


Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.

Predicted band size : 53 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



LPL Antibody (Center) (A00316-1) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of LPL Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Overlay histogram showing HeLa cells stained with A00316-1 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (A00316-1, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was Rabbit IgG (1g/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-LPL Antibody (Center)