

Anti-Survivin BIRC5 Antibody

Catalog Number: A00379-2

About BIRC5

This enzyme is necessary for target cell lysis in cell-mediated immune responses. It cleaves after Asp. Seems to be linked to an activation cascade of caspases (aspartate-specific cysteine proteases) responsible for apoptosis execution. Cleaves caspase-3, -7, -9 and 10 to give rise to active enzymes mediating apoptosis.

David A. Jans, J. Biol. Chem., Nov 1996; 271: 30781. Eugene Estella, Diabetes, Aug 2006; 55: 2212 - 2219. Yong Gu, J. Biol. Chem., May 1996; 271: 10816. Jiuru Sun, J. Biol. Chem., Nov 1996; 271: 27802.

Overview

Product Name	Anti-Survivin BIRC5 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Survivin BIRC5 Antibody catalog # A00379-2. Tested in WB,ICC/IF,IHC,Flow Cytometry applications. This antibody reacts with Human.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O15392

Technical Details

Immunogen	Synthesized peptide derived from human CREB-2 around the phosphorylation site of S219.
Predicted Reactive Species	Canine, Monkey
Cross Reactivity	Does not cross-react with eNOS or nNOS.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml





888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	ProA affinity purified
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:1,000 ICC: 1:50-1:200 IHC: 1:50-1:200 FC: 1:50-1:100



Anti-Survivin BIRC5 Antibody (A00379-2) Images




Figure 1. Immunocytochemistry staining of BIRC5 using Anti-Survivin BIRC5 Antibody (A00379-2).

ICC staining Survivin in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde




Figure 2. Immunohistochemistry validation of BIRC5 using Anti-Survivin BIRC5 Antibody (A00379-2).

Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Survivin antibody. Counter stained with hematoxylin.

For more protocol information of IHC

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Survivin BIRC5 Antibody