

Anti-ULK1 (phospho-Ser636) Antibody

Catalog Number: A00584S636

About ULK1

Transcriptional regulator that binds to DNA as a dimer or as a tetramer, but not as a monomer. Binds to G-doublets in an A/T-rich environment; the preferred motif is a tandem repeat of 5'-ATTGGTTA-3' combined with a 5'-TTATTA-3' box. Binds to nucleosomes By similarity. Binds to chromatin and interacts selectively with histone H3 that is not methylated at 'Lys-4', not phosphorylated at 'Thr-3' and not methylated at 'Arg-2'. Functions as a sensor of histone H3 modifications that are important for the epigenetic regulation of gene expression. Functions as a transcriptional activator and promotes the expression of otherwise tissue-specific self-antigens in the thymus, which is important for self tolerance and the avoidance of autoimmune reactions.

Nagamine K., Nat. Genet. 17:393-398(1997).

Aaltonen J., Nat. Genet. 17:399-403(1997).

Hattori M., Nature 405:311-319(2000).

Overview

Product Name	Anti-ULK1 (phospho-Ser636) Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ULK1 (phospho-Ser636) Antibody catalog # A00584S636. Tested in WB applications. This antibody reacts with Human,Mouse,Rat.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O75385

Technical Details

Immunogen	Synthesized peptide derived from human EphA2/5 around the phosphorylation site of Y594.
Predicted Reactive Species	Canine, Monkey
Isotype	IgG
Form	Liquid

Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1000

Anti-ULK1 (phospho-Ser636) Antibody (A00584S636) Images




Figure 1. Western blotting validation for Anti-ULK1 (phospho-Ser636) Antibody A00584S636

Western blot (WB) analysis of ULK1 (Phospho-Ser638) polyclonal antibody at 1:500 dilution
Lane1:L02 whole cell lysate
Lane2:L02 treated with PBS(1

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ULK1 (phospho-Ser636) Antibody