

Anti-Prolactin/PRL Antibody Picoband™

Catalog Number: A00601-1

About PRL

Prolactin (PRL) also known as luteotropic hormone (LTH) is a protein that in humans is encoded by the PRL gene. Prolactin is a peptide hormone discovered by Henry Friesen. Although it is perhaps best known for its role in lactation, prolactin already existed in the oldest known vertebrates—fishes—where its most important functions were probably related to control of water and salt balance. Prolactin also acts in a cytokine-like manner and as an important regulator of the immune system. Prolactin has important cell cycle related functions as a growth-, differentiating- and anti-apoptotic factor. As a growth factor binding to cytokine like receptors it has also profound influence on hematopoiesis, angiogenesis and is involved in the regulation of blood clotting through several pathways.

Overview

Product Name	Anti-Prolactin/PRL Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Prolactin/PRL Antibody Picoband™ catalog # A00601-1. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P01236

Technical Details

Immunogen	E.coli-derived human Prolactin/PRL recombinant protein (Position: A36-D206).
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5 µg/ml, Mouse, Rat Direct ELISA, 0.1-0.5 µg/ml, Human

Anti-Prolactin/PRL Antibody Picoband™ (A00601-1) Images




Figure 1. Western blot analysis of Prolactin/PRL using anti-Prolactin/PRL antibody (A00601-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.
Lane 1: rat brain tissue lysates,
Lane 2: mouse brain tissue lysates.
After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Prolactin/PRL antigen affinity purified polyclonal antibody (Catalog # A00601-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Prolactin/PRL at approximately 26 kDa. The expected band size for Prolactin/PRL is at 26 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Prolactin/PRL Antibody