

Anti-CXCR5 Antibody Picoband™

Catalog Number: A00663-2

About CXCR5

CXCR5(Chemokine CXC Motif Receptor 5), also known as BLR1, is a G protein-coupled seven transmembrane receptor for chemokine CXCL13(also known as BLC) and belongs to the CXC chemokine receptor family. In humans, the CXC-R5 protein is encoded by the CXCR5 gene. The gene plays an essential role in B cell migration. Reif et al.(2002) concluded that their findings defined the mechanism of B-cell relocalization in response to antigen, and established that cell position in vivo can be determined by the balance of responsiveness to chemoattractants made in separate but adjacent zones. Chan et al.(2003) investigated the expression of chemokines and chemokine receptors in eyes with primary intraocular B-cell lymphoma(PIOL).

Overview

Product Name	Anti-CXCR5 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CXCR5 Antibody Picoband™ catalog # A00663-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P32302

Technical Details

Immunogen	E.coli-derived human CXCR5 recombinant protein (Position: A181-S358).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Flow Cytometry, 1-3 ug/1x10⁶ cells, Human Direct ELISA, 0.1-0.5 ug/ml, Human

Anti-CXCR5 Antibody Picoband™ (A00663-2) Images




Figure 1. Western blot analysis of CXCR5 using anti-CXCR5 antibody (A00663-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Raji whole cell lysates,
Lane 2: human MOLT-4 whole cell lysates,
Lane 3: human Hela whole cell lysates,
Lane 4: rat brain tissue lysates,
Lane 5: mouse brain tissue lysates,
Lane 6: mouse lung tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CXCR5 antigen affinity purified polyclonal antibody (Catalog # A00663-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CXCR5 at approximately 50 kDa. The expected band size for CXCR5 is at 42 kDa.




Figure 2. Flow Cytometry analysis of HEL cells using anti-CXCR5 antibody (A00663-2).

Overlay histogram showing HEL cells stained with A00663-2 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CXCR5 Antibody (A00663-2, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CXCR5 Antibody