

Anti-PKC Delta (phospho-S645) PRKCD Antibody

Catalog Number: A00822S645

About PRKCD

Collagen type III occurs in most soft connective tissues along with type I collagen. Involved in regulation of cortical development. Is the major ligand of GPR56 in the developing brain and binding to GPR56 inhibits neuronal migration and activates the RhoA pathway by coupling GPR56 to GNA13 and possibly GNA12.

Arja Lamberg, J. Biol. Chem., May 1996; 271: 11988.

GW Conrad, J. Cell Biol., Mar 1980; 84: 501.

DR Keene, J. Cell Biol., Nov 1987; 105: 2393.

HM Wang, J. Histochem. Cytochem., Nov 1980; 28: 1215.

Overview

Product Name	Anti-PKC Delta (phospho-S645) PRKCD Antibody
Reactive Species	Human
Description	Boster Bio Anti-PKC Delta (phospho-S645) PRKCD Antibody catalog # A00822S645. Tested in WB,IHC applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q05655

Technical Details

Immunogen	Synthesized peptide derived from human CYP21A2
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1000 IHC: 1:50-1:200

Anti-PKC Delta (phospho-S645) PRKCD Antibody (A00822S645) Images




Figure 2. Immunohistochemistry validation of PRKCD using Anti-PKC Delta (phospho-S645) PRKCD Antibody (A00822S645).

Immunohistochemistry (IHC) analyzes of p-PKC delta (S645) pAb in paraffin-embedded human breast carcinoma tissue at 1:50.

For more protocol information of IHC




Figure 1. Western blotting validation for Anti-PKC Delta (phospho-S645) PRKCD Antibody A00822S645

Western blot (WB) analysis of p-PKC delta (S645) pAb at 1:500 dilution

Lane1:A549 whole cell lysate(20ug)

Lane2:HCT116 whole cell lysate(20ug)

Lane3:SGC7901 whole cell lysate(40ug)

Lane4:The Uterus tissue lysate of Mouse(40ug)

Lane5:The Uterus tissue lysate of Rat(40ug)

Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PKC Delta (phospho-S645) PRKCD Antibody