

Anti-NEDD8 LRP1 Antibody

Catalog Number: A00829-1

About LRP1

Collagen type III occurs in most soft connective tissues along with type I collagen. Involved in regulation of cortical development. Is the major ligand of GPR56 in the developing brain and binding to GPR56 inhibits neuronal migration and activates the RhoA pathway by coupling GPR56 to GNA13 and possibly GNA12.

Arja Lamberg, J. Biol. Chem., May 1996; 271: 11988.

GW Conrad, J. Cell Biol., Mar 1980; 84: 501.

DR Keene, J. Cell Biol., Nov 1987; 105: 2393.

HM Wang, J. Histochem. Cytochem., Nov 1980; 28: 1215.

Overview

Product Name	Anti-NEDD8 LRP1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-NEDD8 LRP1 Antibody catalog # A00829-1. Tested in WB,ICC/IF,IHC,IP applications. This antibody reacts with Human,Mouse,Rat.
Application	IP, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q07954

Technical Details

Immunogen	Synthesized peptide derived from human TRAP230
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	ProA affinity purified
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1,000 ICC: 1:50-1:200 IHC: 1:50-1:100 IP: 1:50

Anti-NEDD8 LRP1 Antibody (A00829-1) Images




Figure 1. Immunocytochemistry staining of LRP1 using Anti-NEDD8 LRP1 Antibody (A00829-1).

ICC staining NEDD8 in HepG2 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde




Figure 2. Immunohistochemistry validation of LRP1 using Anti-NEDD8 LRP1 Antibody (A00829-1).

Immunohistochemical analysis of paraffin-embedded human lung cancer tissue using anti-NEDD8 antibody. Counter stained with hematoxylin.

For more protocol information of IHC

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NEDD8 LRP1 Antibody