

Anti-Bone marrow stromal antigen 2 Bst2 Antibody

Catalog Number: A00934

About BST2

Bst2 was first identified as a novel terminal B-cell-restricted antigen termed HM1.24 that is expressed on certain bone marrow cells as well as other tissues. The surface expression of Bst2 on fibroblast cell lines facilitated the stromal cell-dependent growth of DW34, a pre-B-cell line, suggesting that this protein is involved in pre-B-cell growth. Later experiments also showed it to be upregulated in several myeloma cells. More recently, Bst2 was identified an inhibitor of retroviral release from human cells whose activity is antagonized by the HIV-1 accessory protein Vpu. While Bst2 caused retention of virions on cell surfaces or endocytosis into Bst2-positive compartments, its depletion abolished the viral requirement for Vpu for virus release. This activity may represent a potential therapeutic strategy for the treatment of HIV/AIDS.

Overview


Product Name	Anti-Bone marrow stromal antigen 2 Bst2 Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Bone marrow stromal antigen 2 Bst2 Antibody (Catalog # A00934). Tested in ELISA, WB applications. This antibody reacts with Human, Mouse.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Bst2 Antibody is supplied in PBS containing 0.02% sodium azide.
Storage Instructions	Bst2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Host	Rabbit
Uniprot ID	Q8R2Q8

Technical Details

Immunogen	Bst2 antibody was raised against a 18 amino acid synthetic peptide near the amino terminus of the mouse Bst2. The immunogen is located within amino acids 60 - 110 of Bst2.
Predicted Reactive Species	Rat
Cross Reactivity	ARF6 antibody is human, mouse and rat reactive.
Isotype	IgG
Form	Liquid
Concentration	1 mg/mL

Purification	Bst2 Antibody is affinity chromatography purified via peptide column.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Bst2 antibody can be used for detection of Bst2 by Western blot at 1 - 2 ug/mL. Antibody validated: Western Blot in human samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.

Anti-Bone marrow stromal antigen 2 Bst2 Antibody (A00934) Images



Western blot analysis of Bst2 in Daudi cell lysate with Bst2 antibody at (A) 1 and (B) 2 ug/mL.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Bone marrow stromal antigen 2 Bst2 Antibody