

# **Anti-Tal1 Antibody Picoband™**

Catalog Number: A00944-2

#### **About TAL1**

T-cell acute lymphocytic leukemia protein 1 is a protein that in humans is encoded by the TAL1 gene. The protein encoded by TAL1 is a basic helix-loop-helix transcription factor. TAL1 is a transcriptionally complex gene that is expressed throughout development, activating or repressing transcription in hematopoietic, neural, and endothelial precursors. This gene is mapped to chromosome 1p33.

#### Overview

| Product Name         | Anti-Tal1 Antibody Picoband™  |
|----------------------|---|
| Reactive Species     | Human, Mouse, Rat   |
| Description          | Boster Bio Anti-Tal1 Antibody Picoband™ catalog # A00944-2. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.  |
| Application          | ELISA, WB   |
| Clonality            | Polyclonal  |
| Formulation          | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .  |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit  |
| Uniprot ID           | P17542  |

#### **Technical Details**

| Immunogen                     | E. coli-derived human Tal1 recombinant protein (Position: Y174-D259).                           |
|-------------------------------|---|
| Predicted Reactive Species    | Human   |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity              | No cross-reactivity with other proteins.  |
| Isotype                       | Rabbit IgG  |
| Form                          | Lyophilized   |
| Concentration                 | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.                       |
| Purification                  | Immunogen affinity purified.  |
|                               |   |



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blot, 0.1-0.5ug/ml  Direct ELISA, 0.1-0.5ug/ml |
|---------------------|--|
|---------------------|--|



## Anti-Tal1 Antibody Picoband™ (A00944-2) Images




Figure 1. Western blot analysis of Tal1 using anti-Tal1 antibody (A00944-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human U20S cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Tal1 antigen affinity purified polyclonal antibody (Catalog # A00944-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Tal1 at approximately 39KD. The expected band size for Tal1 is at 34KD.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Tal1 Antibody ™