

Anti-NeuroD1 Antibody

Catalog Number: A01038

About NEUROD1

4E-BP1 encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.

Gingras AC, et al. (1998) Genes Dev 12(4): 502-513.

Brugarolas J, et al. (2004) Genes Dev 18(23): 2893-2904.

Kumar V, et al. (2000) EMBO J 19(5): 1087-1097.

Moody CA, et al. (2005) J Virol 79(9): 5499-5506.

Burnett PE, et al. (1998) Proc Natl Acad Sci U S A 95(4): 1432-1437.

Overview

| Product Name | Anti-NeuroD1 Antibody |
|----------------------|--|
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-Neuro D1 Antibody catalog # A01038. Tested in WB,IP applications. This antibody reacts with Human,Mouse,Rat. |
| Application | IP, WB |
| Clonality | Polyclonal |
| Formulation | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2 |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q13562 |

Technical Details

| Immunogen | Synthesized peptide derived from human Paxillin around the phosphorylation site of Y88. |
|----------------------------|---|
| Predicted Reactive Species | Boar, Bovine, Canine, Golden Hamster |
| Cross Reactivity | No cross reactivity with other proteins. |
| Isotype | IgG |
| | |





888-466-3604 | support@bosterbio.com | www.bosterbio.com

| Form | Liquid |
|---------------------|--|
| Concentration | 1 mg/ml |
| Purification | ProA affinity purified |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1,000 IP: 1:10-1:50 |



Anti-NeuroD1 Antibody (A01038) Images




Figure 1. Western blotting validation for Anti-NeuroD1 Antibody A01038

Western blot analysis of NeuroD1 on different cells lysates using anti-NeuroD1 antibody at 1/500 dilution. Positive control: Line 1: human brain Line 2:SH-SY5Y Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-NeuroD1 Antibody