

Anti-Scavenging Receptor SR-BI/SCARB1 Antibody Picoband™

Catalog Number: A01093-1

About SCARB1

Scavenger receptor class B member 1 (SRB1), also known as SR-BI, is a protein that in humans is encoded by the SCARB1 gene. SR-BI functions as a receptor for high-density lipoprotein. Scavenger receptor class B, type I (SR-BI) is an integral membrane protein found in numerous cell types/tissues, including the liver and adrenal. It is best known for its role in facilitating the uptake of cholesteryl esters from high-density lipoproteins in the liver. This process drives the movement of cholesterol from peripheral tissues towards the liver for excretion. This movement of cholesterol is known as reverse cholesterol transport and is a protective mechanism against the development of atherosclerosis, which is the principal cause of heart disease and stroke. SR-BI has also been identified in the livers of non-mammalian species (turtle, goldfish, shark, chicken, frog, and skate), suggesting it emerged early in vertebrate evolutionary history. The turtle also seems to upregulate SB-RI during egg development, indicating that cholesterol efflux may be at peak levels during developmental stages.

Overview

Product Name	Anti-Scavenging Receptor SR-BI/SCARB1 Antibody Picoband™
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Scavenging Receptor SR-BI/SCARB1 Antibody Picoband™ catalog # A01093-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q8WTV0

Technical Details

Immunogen	E.coli-derived human Scavenging Receptor SR-BI/SCARB1 recombinant protein (Position: F70-R492).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5 ug/ml, Human, Mouse Flow Cytometry, 1-3 ug/1x10⁶ cells, Human Direct ELISA, 0.1-0.5 ug/ml, Human

Anti-Scavenging Receptor SR-BI/SCARB1 Antibody Picoband™ (A01093-1) Images




Figure 1. Western blot analysis of Scavenging Receptor SR-BI/SCARB1 using anti-Scavenging Receptor SR-BI/SCARB1 antibody (A01093-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human T-47D whole cell lysates,

Lane 2: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Scavenging Receptor SR-BI/SCARB1 antigen affinity purified polyclonal antibody (Catalog # A01093-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Scavenging Receptor SR-BI/SCARB1 at approximately 85 kDa. The expected band size for Scavenging Receptor SR-BI/SCARB1 is at 85 kDa.




Figure 2. Flow Cytometry analysis of MCF-7 cells using anti-Scavenging Receptor SR-BI/SCARB1 antibody (A01093-1).

Overlay histogram showing MCF-7 cells stained with A01093-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Scavenging Receptor SR-BI/SCARB1 Antibody (A01093-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Scavenging Receptor SR-BI/SCARB1 Antibody