

Anti-TAPA1/Cd81 Antibody Picoband™

Catalog Number: A01281-3

About Cd81

CD81 molecule, also known as CD81 (Cluster of Differentiation 81), is a protein which in humans is encoded by the CD81 gene. The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. This protein appears to promote muscle cell fusion and support myotube maintenance. Also it may be involved in signal transduction. This gene is localized in the tumor-suppressor gene region and thus it is a candidate gene for malignancies. Two transcript variants encoding different isoforms have been found for this gene.

Overview

Product Name	Anti-TAPA1/Cd81 Antibody Picoband™
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-TAPA1/Cd81 Antibody Picoband™ catalog # A01281-3. Tested in ELISA, IF, IHC, WB applications. This antibody reacts with Mouse, Rat.
Application	ELISA, IF, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P35762

Technical Details

Immunogen	E. coli-derived mouse TAPA1 recombinant protein (Position: K116-K201).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml Immunofluorescence, 5ug/ml Direct ELISA, 0.1-0.5ug/ml

Anti-TAPA1/Cd81 Antibody Picoband™ (A01281-3) Images




Figure 1. Western blot analysis of TAPA1/CD81 using anti-TAPA1/CD81 antibody (A01281-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,
Lane 2: mouse ovary tissue lysates,
Lane 3: mouse brain tissue lysates,
Lane 4: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TAPA1/CD81 antigen affinity purified polyclonal antibody (Catalog # A01281-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TAPA1/CD81 at approximately 22 kDa. The expected band size for TAPA1/CD81 is at 26 kDa.




Figure 2. IHC analysis of TAPA1/CD81 using anti-TAPA1/CD81 antibody (A01281-3).

TAPA1/CD81 was detected in a paraffin-embedded section of rat liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TAPA1/CD81 Antibody (A01281-3) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 3. IF analysis of TAPA1/CD81 using anti-TAPA1/CD81 antibody (A01281-3).

TAPA1/CD81 was detected in a paraffin-embedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-TAPA1/CD81 Antibody (A01281-3) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 4. IF analysis of TAPA1/CD81 using anti-TAPA1/CD81 antibody (A01281-3). TAPA1/CD81 was detected in a paraffin-embedded section of rat liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-TAPA1/CD81 Antibody (A01281-3) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

2 Publications Citing This Product

1. PubMed ID: 32059163, Wang D,Hao C,Zhang L,Zhang J,Liu S,Li Y,Qu Y,Zhao Y,Huang R,Wei J,Yao W.Exosomal miR-125a-5p derived from silica-exposed macrophages induces fibroblast transdifferentiation.Ecotoxicol Environ Saf.2020 Apr 1;192:110253.doi:10.1016/j.ecoenv.2020.110253.Epub
2. PubMed ID: 32400849, Cao G,Meng X, Han X,Li J.Exosomes derived from circRNA Rtn4-modified BMSCs attenuate TNF-alpha-induced cytotoxicity and apoptosis in murine MC3T3-E1 cells by sponging miR-146a.Biosci Rep.2020 May 29;40(5): BSR20193436.doi:10.1042/BSR20193436.PMID:32400849;PMC

Visit bosterbio.com/anti-tapa1-picoband-trade-antibody-a01281-3-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TAPA1/Cd81 Antibody [™]