

Anti-ENPP1 Antibody

Catalog Number: A01342

About ENPP1

The NCOA3 (also known as nuclear receptor coactivator 3 isoform a) is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions in a hormone-dependent fashion. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. NCOA3 probably acts via remodeling of chromatin and is involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). NCOA3 is also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit. NCOA3 may be associated with diseases such as breast cancer and meningothelial meningioma. Anti-NCOA3 Antibody is useful for researchers interested in cancer research, chromatin binding, transcription coactivator activity, Immunology, and Nuclear Signaling research.

Overview

Product Name	Anti-ENPP1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ENPP1 Antibody catalog # A01342. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P22413

Technical Details

Immunogen	Synthesized peptide derived from human NCoA-3.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	ENPP1 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB 1:500-2000 ELISA 1:5000-20000

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ENPP1 Antibody