

# Anti-Trk B (phospho-Y515) NTRK2 Antibody

Catalog Number: A01388Y515

#### **About NTRK2**

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection.

Abujiang Pataer, et,al. (2002) Cancer Res; 62: 2239. K. D. Ryman, et,al. (2005) J. Virol; 79: 1487 - 1499.

Susana Guerra, et,al. (2006) J. Biol. Chem; 281: 18734 - 18745.

### Overview

Product Name	Anti-Trk B (phospho-Y515) NTRK2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Trk B (phospho-Y515) NTRK2 Antibody catalog # A01388Y515. Tested in WB,IHC applications. This antibody reacts with Human,Mouse,Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16620

## **Technical Details**

Immunogen	Synthesized peptide derived from human Emt around the phosphorylation site of Y512.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml



888-466-3604 | support@bosterbio.com | www.bosterbio.com



Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitopespecific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  WB: 1:500-1:1000  IHC: 1:50-1:200



# Anti-Trk B (phospho-Y515) NTRK2 Antibody (A01388Y515) Images




Figure 1. Immunohistochemistry validation of NTRK2 using Anti-Trk B (phospho-Y515) NTRK2 Antibody (A01388Y515).

Immunohistochemistry (IHC) analyzes of p-Trk B (Y515) pAb in paraffin-embedded human colorectal carcinoma tissue at 1:50.

For more protocol information of IHC

# Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Trk B (phospho-Y515) NTRK2 Antibody