

Anti-TrkB (Phospho-Y817) NTRK2 Antibody

Catalog Number: A01388Y817

About NTRK2

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection.

Abujiang Pataer, et,al. (2002) Cancer Res; 62: 2239. K. D. Ryman, et,al. (2005) J. Virol; 79: 1487 - 1499.

Susana Guerra, et,al. (2006) J. Biol. Chem; 281: 18734 - 18745.

Overview

Product Name	Anti-TrkB (Phospho-Y817) NTRK2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-TrkB (Phospho-Y817) NTRK2 Antibody catalog # A01388Y817. Tested in WB,ICC/IF,IHC,IP applications. This antibody reacts with Human,Mouse,Rat.
Application	IP, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16620

Technical Details

Immunogen	Synthesized peptide derived from human Trk B around the phosphorylation site of Y706.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml





BOSTER
antibody and ELISA experts

Purification	ProA affinity purified
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:1,000-1:5,000 ICC: 1:50-1:200 IHC: 1:50-1:200



Anti-TrkB (Phospho-Y817) NTRK2 Antibody (A01388Y817) Images




Figure 2. Immunocytochemistry staining of NTRK2 using Anti-TrkB (Phospho-Y817) NTRK2 Antibody (A01388Y817).

ICC staining phospho -TrkB(Y817) in PANC-1 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde




Figure 1. Western blotting validation for Anti-TrkB (Phospho-Y817) NTRK2 Antibody A01388Y817

Western blot analysis of phospho -TrkB(Y817) on mouse brain lysates using anti-phospho -TrkB(Y817) antibody at 1/1

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-TrkB (Phospho-Y817) NTRK2 Antibody