

# Anti-BNIP-3 (E96) Antibody

Catalog Number: A01469-1

### **About BNIP3**

Type I receptor for TGF-beta family ligands BMP9/GDF2 and BMP10 and important regulator of normal blood vessel development. On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. May bind activin as well.

ten Dijke P., Oncogene 8:2879-2887(1993). Attisano L., Cell 75:671-680(1993). Berg J.N., Am. J. Hum. Genet. 61:60-67(1997).

### Overview

| Product Name         | Anti-BNIP-3 (E96) Antibody   |
|----------------------|--|
| Reactive Species     | Human, Mouse, Rat  |
| Description          | Boster Bio Anti-BNIP-3 (E96) Antibody catalog # A01469-1. Tested in IHC applications. This antibody reacts with Human, Mouse, Rat.         |
| Application          | IHC  |
| Clonality            | Polyclonal   |
| Formulation          | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2   |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit   |
| Uniprot ID           | Q12983   |

### **Technical Details**

| Immunogen                  | Synthesized peptide derived from internal of human ACVL1. |
|----------------------------|---|
| Predicted Reactive Species | Boar, Bovine, Canine, Golden Hamster                      |
| Cross Reactivity           | No cross reactivity with other proteins.                  |
| Isotype                    | IgG   |
| Form                       | Liquid  |
| Concentration              | 1 mg/ml   |



888-466-3604 | support@bosterbio.com | www.bosterbio.com

| Purification        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).   |
|---------------------|---|
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  IHC: 1:50-1:200 |

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-BNIP-3 (E96) Antibody