

Anti-RAC2 Antibody

Catalog Number: A01714

About RAC2

Regulates and initiates biosynthesis of N-acetylneuraminic acid (NeuAc), a precursor of sialic acids. Plays an essential role in early development By similarity. Required for normal sialylation in hematopoietic cells. Sialylation is implicated in cell adhesion, signal transduction, tumorigenicity and metastatic behavior of malignant cells.

Lucka L., FEBS Lett. 454:341-344(1999).

Seppala R., Am. J. Hum. Genet. 64:1563-1569(1999).

Huizing M., Submitted (OCT-2000) to the EMBL/GenBank/DDBJ databases.

Overview

Product Name	Anti-RAC2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-RAC2 Antibody catalog # A01714. Tested in WB applications. This antibody reacts with Human,Mouse,Rat.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P15153

Technical Details

Immunogen	Synthesized peptide derived from human GRIP-1 around the phosphorylation site of S736.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1000

Anti-RAC2 Antibody (A01714) Images




Figure 1. Western blotting validation for Anti-RAC2 Antibody A01714

Western blot (WB) analysis of RAC2 polyclonal antibody at 1:500 dilution

Lane1: Jurkat whole cell lysate

Lane2: sp2/0 whole cell lysate

Lane3: H9C2 whole cell lysate

Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RAC2 Antibody