

Anti-Phospho-PKC beta (S661) PRKCB Antibody

Catalog Number: A01940S661

About PRKCB

Cullins assemble a potentially large number of ubiquitin ligases by binding to the RING protein ROC1 to catalyse polyubiquitination, as well as binding to various specificity factors to recruit substrates. Cullin 5 is a component of E3 ubiquitin ligase complexes, which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. Cullin 5 seems to be involved proteasomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa protein. Cullin 5 may form a cell surface vasopressin receptor. Cullin 5 is part of a E3 ubiquitin ligase complex with elongin BC complex (TCEB1 and TCEB2), RBX1 and MUF1, complexes with elongin BC complex (TCEB1 and TCEB2), RXB1 and TCEB3 or SOCS1 or WSB1; elongin BC complex (TCEB1 and TCEB2), RBX1 and VHL; elongin BC complex (TCEB1 and TCEB2), RBX1, adenovirus type 5 E1B-55kDa protein and adenovirus type 5 E4-orf6. Cullin 5 Interacts with RBX1 and RNF7.

Overview

Product Name	Anti-Phospho-PKC beta (S661) PRKCB Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Phospho-PKC beta (S661) PRKCB Antibody catalog # A01940S661. Tested in ELISA, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05771

Technical Details

Immunogen	Synthesized peptide derived from human PKC beta around the phosphorylation site of S661.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Phospho-PKC beta (S661) PRKCB Antibody