

Anti-GPBAR1 Antibody

Catalog Number: A01958

About GPBAR1

Cullins assemble a potentially large number of ubiquitin ligases by binding to the RING protein ROC1 to catalyse polyubiquitination, as well as binding to various specificity factors to recruit substrates. Cullin 5 is a component of E3 ubiquitin ligase complexes, which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. Cullin 5 seems to be involved proteasomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa protein. Cullin 5 may form a cell surface vasopressin receptor. Cullin 5 is part of a E3 ubiquitin ligase complex with elongin BC complex (TCEB1 and TCEB2), RBX1 and MUF1, complexes with elongin BC complex (TCEB1 and TCEB2), RXB1 and TCEB3 or SOCS1 or WSB1; elongin BC complex (TCEB1 and TCEB2), RBX1 and VHL; elongin BC complex (TCEB1 and TCEB2), RBX1, adenovirus type 5 E1B-55kDa protein and adenovirus type 5 E4-orf6. Cullin 5 Interacts with RBX1 and RNF7.

Overview

Product Name	Anti-GPBAR1 Antibody
Reactive Species	Human
Description	Boster Bio Anti-GPBAR1 Antibody catalog # A01958. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8TDU6

Technical Details

Immunogen	Synthesized peptide derived from human IBP5 protein.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

WB: 1:500-1:1000

Anti-GPBAR1 Antibody (A01958) Images




Figure 1. Western blotting validation for Anti-GPBAR1 Antibody A01958

Western blot (WB) analysis of GPBAR1 polyclonal antibody at 1:500 dilution

Lane1:HEK293T whole cell lysate

Lane2:Raw264.7 whole cell lysate

Lane3:PC12 whole cell lysate

Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-GPBAR1 Antibody