

Anti-PERK EIF2AK3 Antibody

Catalog Number: A01992

About EIF2AK3

The PKR-like endoplasmic reticulum kinase (PERK, also known as Eukaryotic translation initiation factor 2- α kinase 3) is a type I transmembrane protein localized to the endoplasmic reticulum (ER). PERK consists of an N-terminal ER luminal domain, a membrane-spanning region, and a cytosolic C-terminal serine/threonine kinase domain (1). The luminal domain of PERK is bound to the ER chaperone GRP78 in unstressed cells (2). PERK activation occurs upon accumulation of misfolded proteins in the ER lumen, which triggers GRP78 dissociation from PERK thereby allowing PERK dimerization and autophosphorylation (3, 4). PERK phosphorylates two established targets: the eukaryotic translation initiation factor 2 α (eIF2, (1)) and the Nrf2 transcription factor (5). Phosphorylation of eIF2 results in attenuation of translation initiation (6). The translational block also contributes to cell cycle arrest due to loss of the G1 regulatory protein, cyclin D1 (7). PERK-dependent phosphorylation of Nrf2 promotes transcription of phase II detoxifying enzymes which is critically important for elimination of intracellular reactive oxygen species (8). Thus, while inhibiting new protein synthesis and thereby decreasing the ER protein load PERK simultaneously induces expression of genes that help restore cellular redox homeostasis and promote survival.

Overview


Product Name	Anti-PERK EIF2AK3 Antibody
Reactive Species	Mouse
Description	Boster Bio Anti-PERK EIF2AK3 Antibody (Catalog # A01992). Tested in IHC, IP, WB applications. This antibody reacts with Mouse.
Application	IP, IHC, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	Q9Z2B5

Technical Details


Immunogen	This whole rabbit serum was prepared by repeated immunizations with a recombinant fusion protein from amino acids 601-1115 of mouse deltaN PERK.
Predicted Reactive Species	Bovine, Canine, Equine, Guinea Pig, Rabbit

Isotype	Antiserum
Form	Liquid (sterile filtered)
Concentration	80 mg/mL by Refractometry
Purification	This antiserum is directed against PERK and reacts with the PERK from mouse tissues. Reactivity to other species is unknown.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>ELISA: 1:4,000 - 1:20,000</p> <p>IHC: 1:1,000</p> <p>IF Microscopy: User optimized</p> <p>IP: 10-30ul</p> <p>WB: 1:500 - 1:3000</p>

Anti-PERK EIF2AK3 Antibody (A01992) Images



Western blot analysis of PERK expression in cell lysates. 300ug PERK over-expressing 293T cell lysate (lanes 1 & 2), or 800ug wild type (Lanes 3 & 4), and PERK knock out (lanes 5 & 6) MEF cell lysate were immunoprecipitated with 15ul anti-PERK. PERK was detected using rabbit anti-PERK Antiserum, not purified polyclonal antibody (Catalog # A01992. Lane 1, 293T cells over-expressing Myc-PERK wt, Lane 2, 293T cells over-expressing Myc-PERK K618A. Personal Communication. A, Diehl, Univ. of Pennsylvania, Philadelphia, PA.



PERK was detected in paraffin-embedded sections of mouse mammary gland tissues using rabbit anti-PERK Antiserum, not purified polyclonal antibody (Catalog # A01992) at 1:1000. Positive staining signal observed in wild type mouse sample with anti-PERK staining only (middle image), but not in the knock out mouse sample (right image) and pre-immune serum staining (left image) The anti-PERK was diluted 1:1,000 in 5% goat serum in PBS and allowed to incubate for 2h at room temperature in a humidified chamber. Personal Communication. A, Diehl, Univ. of Pennsylvania, Philadelphia, PA.

2 Publications Citing This Product

1. PubMed ID: 10.1038/s41419-020-02930-y, BCAT1 binds the RNA-binding protein ZNF423 to activate autophagy via the IRE1-XBP-1-RIDD axis in hypoxic PSMCs
2. PubMed ID: 27293989, Recombinant Newcastle disease virus (rL-RVG) triggers autophagy and apoptosis in gastric carcinoma cells by inducing ER stress

Visit bosterbio.com/anti-perk-antibody-a01992-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PERK EIF2AK3 Antibody