

Anti-TRPC4 Antibody Picoband™

Catalog Number: A02037-1

About TRPC4

The short transient receptor potential channel 4 (TRPC4), also known as Trp-related protein 4, is a protein that in humans is encoded by the TRPC4 gene. TRPC4 is a member of the transient receptor potential cation channels. It is mapped to 13q13.3. This protein forms a non-selective calcium-permeable cation channel that is activated by Gq-coupled receptors and tyrosine kinases, and plays a role in multiple processes including endothelial permeability, vasodilation, neurotransmitter release and cell proliferation. TRPC4 also has an essential role in endothelial-dependent regulation of vascular tone, endothelial permeability, and neurotransmitter release from thalamic interneurons.

Overview

Product Name	Anti-TRPC4 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-TRPC4 Antibody Picoband™ catalog # A02037-1. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UBN4

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human TRPC4, identical to the related mouse and rat sequences.
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml

Anti-TRPC4 Antibody Picoband™ (A02037-1) Images




Figure 1. Western blot analysis of TRPC4 using anti-TRPC4 antibody (A02037-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human COLO-320 whole cell lysates,

Lane 3: human MCF-7 whole cell lysates,

Lane 4: human 293T whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes.

Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TRPC4 antigen affinity purified polyclonal antibody (Catalog # A02037-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TRPC4 at approximately 150KD. The expected band size for TRPC4 is at 112KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TRPC4 Antibody™