

Anti-Fatty Acid Binding Protein 5/FABP5 Antibody Picoband™

Catalog Number: A02083

About Fabp5

FABP5, Fatty acid-binding protein, epidermal, is a protein that in humans is encoded by the FABP5 gene. This gene encodes the fatty acid binding protein found in epidermal cells, and was first identified as being upregulated in psoriasis tissue. It is mapped to 8q21.13. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. It is thought that FABPs roles include fatty acid uptake, transport, and metabolism.

Overview

Product Name	Anti-Fatty Acid Binding Protein 5/FABP5 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Fatty Acid Binding Protein 5/FABP5 Antibody Picoband™ catalog # A02083. Tested in IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q05816

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of mouse FABP5, which shares 81.2% amino acid (aa) sequence identity with human FABP5.
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml Immunocytochemistry/Immunofluorescence, 5 ug/ml

Anti-Fatty Acid Binding Protein 5/FABP5 Antibody Picoband™ (A02083) Images




Figure 1. Western blot analysis of FABP5 using anti-FABP5 antibody (A02083).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human RT4 whole cell lysates,

Lane 2: human PC-3 whole cell lysates,

Lane 3: rat lung tissue lysates,

Lane 4: mouse lung tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FABP5 antigen affinity purified polyclonal antibody (Catalog # A02083) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FABP5 at approximately 15 kDa. The expected band size for FABP5 is at 15 kDa.




Figure 2. IF analysis of FABP5 using anti-FABP5 antibody (A02083).

FABP5 was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-FABP5 Antibody (A02083) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

1 Publications Citing This Product

1. PubMed ID: 10.1016/j.wneu.2016.05.063, High FABP5 Versus CRABP II Expression Ratio in Recurrent Craniopharyngiomas: Implications for Future Treatment

Visit bosterbio.com/anti-fabp5-picoband-trade-antibody-a02083-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Fatty Acid Binding Protein 5/FABP5 Antibody [™]