

Anti-PSD95/DLG4 Antibody Picoband™

Catalog Number: A02128

About DLG4

DLG4 discs large homolog 4, also known as PSD95 or SAP-90, is a protein that in humans is encoded by the DLG4 gene. It is a member of the membrane-associated guanylate kinase (MAGUK) family. This gene is mapped to 17p13.1. DLG4 can heteromultimerize with another MAGUK protein, DLG2, and is recruited into NMDA receptor and potassium channel clusters. These two MAGUK proteins may interact at postsynaptic sites to form a multimeric scaffold for the clustering of receptors, ion channels, and associated signaling proteins. Overexpression of DLG4 in hippocampal neurons could drive maturation of glutamatergic synapses. DLG4 can orchestrate synaptic development and it has a role in synapse stabilization and plasticity. Ubiquitination of DLG4 through an MDM2-mediated pathway can regulate AMPA receptor surface expression during synaptic plasticity.

Overview

Product Name	Anti-PSD95/DLG4 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PSD95/DLG4 Antibody Picoband™ catalog # A02128. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P78352

Technical Details

Immunogen	E.coli-derived human PSD95/DLG4 recombinant protein (Position: D18-L711).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5ug/ml, Mouse, Rat Flow Cytometry, 1-3ug/1x10⁶ cells, Human Direct ELISA, 0.1-0.5ug/ml, Human

Anti-PSD95/DLG4 Antibody Picoband™ (A02128) Images




Figure 1. Western blot analysis of PSD95/DLG4 using anti-PSD95/DLG4 antibody (A02128). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.
Lane 1: rat brain tissue lysates,
Lane 2: mouse brain tissue lysates.
After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PSD95/DLG4 antigen affinity purified polyclonal antibody (Catalog # A02128) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PSD95/DLG4 at approximately 99KD. The expected band size for PSD95/DLG4 is at 99KD.




Figure 2. Flow Cytometry analysis of U937 cells using anti-PSD95/DLG4 antibody (A02128). Overlay histogram showing U937 cells stained with A02128 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PSD95/DLG4 Antibody (A02128, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

1 Publications Citing This Product

1. PubMed ID: 10.1016/j.bbrc.2017.06.108, Overexpression cdc42 attenuates isoflurane-induced neurotoxicity in developmental brain of rats

Visit bosterbio.com/anti-psd95-dlg4-picoband-trade-antibody-a02128-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PSD95/DLG4 Antibody