

Anti-FABP2/I-FABP Antibody Picoband™

Catalog Number: A02378

About Fabp2

FABP 2, Fatty acid-binding protein 2, is a protein that in humans is encoded by the FABP2 gene. Using a human cDNA probe, the gene is assigned to chromosome 4 in somatic cell hybrids. FABP 2 gene contains four exons and is an abundant cytosolic protein in small intestine epithelial cells. The FABPs belong to a multigene family with nearly twenty identified members. And FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. Also, they may be responsible in the modulation of cell growth and proliferation.

Overview

Product Name	Anti-FABP2/I-FABP Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-FABP2/I-FABP Antibody Picoband™ catalog # A02378. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P55050

Technical Details

Immunogen	E.coli-derived mouse FABP2/I-FABP recombinant protein (Position: A2-E132). Mouse FABP2/I-FABP shares 77.9% and 92.4% amino acid (aa) sequence identity with human and rat FABP2/I-FABP, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

Anti-FABP2/I-FABP Antibody Picoband™ (A02378) Images




Figure 1. Western blot analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat liver tissue lysates,

Lane 2: rat small intestine tissue lysates,

Lane 3: mouse liver tissue lysates,

Lane 4: mouse small intestine tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FABP2/I-FABP antigen affinity purified polyclonal antibody (Catalog # A02378) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FABP2/I-FABP at approximately 15 kDa. The expected band size for FABP2/I-FABP is at 15 kDa.




Figure 2. IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378).

FABP2/I-FABP was detected in paraffin-embedded section of mouse intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Figure 3. IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378).

FABP2/I-FABP was detected in paraffin-embedded section of rat intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Figure 4. IHC analysis of FABP2/I-FABP using anti-FABP2/I-FABP antibody (A02378).

FABP2/I-FABP was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-FABP2/I-FABP Antibody (A02378) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-FABP2/I-FABP Antibody TM