

Anti-DNA Ligase III/LIG3 Antibody Picoband™

Catalog Number: A02741-2

About LIG3

DNA ligase 3 is an enzyme that, in humans, is encoded by the LIG3 gene. This gene is a member of the DNA ligase family. Each member of this family encodes a protein that catalyzes the joining of DNA ends but they each have a distinct role in DNA metabolism. The protein encoded by this gene is involved in excision repair and is located in both the mitochondria and nucleus, with translation initiation from the upstream start codon allowing for transport to the mitochondria and translation initiation from a downstream start codon allowing for transport to the nucleus. Additionally, alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Overview

Product Name	Anti-DNA Ligase III/LIG3 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-DNA Ligase III/LIG3 Antibody Picoband™ catalog # A02741-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P49916

Technical Details

Immunogen	E.coli-derived human DNA Ligase III/LIG3 recombinant protein (Position: D297-D963).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat

Flow Cytometry, 1-3 ug/ 1×10^6 cells, Human

Direct ELISA, 0.1-0.5 ug/ml, Human

Anti-DNA Ligase III/LIG3 Antibody Picoband™ (A02741-2) Images




Figure 1. Western blot analysis of DNA Ligase III/LIG3 using anti-DNA Ligase III/LIG3 antibody (A02741-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,
Lane 2: human HEK293 whole cell lysates,
Lane 3: human PC-3 whole cell lysates,
Lane 4: human THP-1 whole cell lysates,
Lane 5: rat PC-12 whole cell lysates,
Lane 6: mouse RAW264.7 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DNA Ligase III/LIG3 antigen affinity purified polyclonal antibody (Catalog # A02741-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for DNA Ligase III/LIG3 at approximately 113 kDa. The expected band size for DNA Ligase III/LIG3 is at 113 kDa.




Figure 2. Flow Cytometry analysis of K562 cells using anti-DNA Ligase III/LIG3 antibody (A02741-2). Overlay histogram showing K562 cells stained with A02741-2 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-DNA Ligase III/LIG3 Antibody (A02741-2, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-DNA Ligase III/LIG3 Antibody