

## **Anti-VRL1/TRPV2 Antibody Picoband®**

Catalog Number: A02786-3

#### **About TRPV2**

TRPV2 (Transient Receptor Potential Cation Channel Subfamily V Member 2), also known as VRL1, is a protein that, in humans, is encoded by the TRPV1 gene. The International Radiation Hybrid Mapping Consortium mapped the TRPV2 gene to chromosome 17. This gene encodes an ion channel that is activated by high temperatures above 52C. The protein may be involved in transduction of high-temperature heat responses in sensory ganglia. It is though that in other tissues the channel may be activated by stimuli other than heat.

#### Overview

Product Name	Anti-VRL1/TRPV2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-VRL1/TRPV2 Antibody Picoband® catalog # A02786-3. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9Y5S1

#### **Technical Details**

Immunogen	E.coli-derived human VRL1/TRPV2 recombinant protein (Position: S7-K390).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3ug/1x10 <sup>6</sup> cells, Human ELISA, 0.1-0.5ug/ml, -



#### Anti-VRL1/TRPV2 Antibody Picoband® (A02786-3) Images



Western blot analysis of VRL1/TRPV2 using anti-VRL1/TRPV2 antibody (A02786-3).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: human A549 whole cell lysates,

Lane 2: human A431 whole cell lysates,


Lane 3: rat brain tissue lysates,

Lane 4: mouse brain tissue lysates,

Lane 5: mouse heart tissue lysates,

Lane 6: mouse lung tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VRL1/TRPV2 antigen affinity purified polyclonal antibody (Catalog # A02786-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for VRL1/TRPV2 at approximately 95KD. The expected band size for VRL1/TRPV2 is at 95KD.



Flow Cytometry analysis of HL-60 cells using anti-VRL1/TRPV2 antibody (A02786-3). Overlay histogram showing HL-60 cells stained with A02786-3 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-VRL1/TRPV2 Antibody (A02786-3, 1ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.