

Anti-AVPR1A/V1aR Antibody Picoband™

Catalog Number: A03279

About AVPR1A

Arginine vasopressin receptor 1A (officially called AVPR1A) is one of the three major receptor types for arginine vasopressin (AVPR1B and AVPR2 being the others). It belongs to the subfamily of G-protein coupled receptors which includes AVPR1B, V2R and OXT receptors. This gene is mapped to 12q14.2. AVPR1A is present throughout the brain, as well as in the periphery, in the liver, kidney, and vasculature. The protein encoded by this gene acts as receptor for arginine vasopressin. Its activity is mediated by G proteins which stimulate a phosphatidylinositol-calcium second messenger system. The receptor mediates cell contraction and proliferation, platelet aggregation, release of coagulation factor and glycogenolysis.

Overview

Product Name	Anti-AVPR1A/V1aR Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-AVPR1A/V1aR Antibody Picoband™ catalog # A03279. Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P37288

Technical Details

Immunogen	E.coli-derived human AVPR1A/V1aR recombinant protein (Position: R43-K268).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.25-0.5ug/ml, Human</p> <p>Direct ELISA, 0.1-0.5ug/ml, Human</p>

Anti-AVPR1A/V1aR Antibody Picoband™ (A03279) Images




Figure 1. Western blot analysis of AVPR1A using anti-AVPR1A antibody (A03279).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human A549 whole cell lysates,
Lane 2: human A431 whole cell lysates,
Lane 3: human HEK293 whole cell lysates,
Lane 4: human THP-1 whole cell lysates,
Lane 5: human Caco-2 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-AVPR1A antigen affinity purified polyclonal antibody (Catalog # A03279) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for AVPR1A at approximately 60KD. The expected band size for AVPR1A is at 47KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-AVPR1A/V1aR Antibody™