

## Anti-Glypican-1 Antibody

Catalog Number: A03871

### About GPC1

Glypican-1 antibody is suitable for Cancer, Immunology and Nuclear Signaling research. Glypican-1 (also known as GPC1 or FLJ38078) is a member of the glypican-related integral membrane proteoglycan family (GRIPS). This protein is a heparan sulfate proteoglycan which is composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains and are anchored to the cell surface via a covalent linkage to glycosylphosphatidylinositol (GPI). Glypicans can modify cell signaling pathways and contribute to cellular proliferation and tissue growth. In humans, glypican-1 is over expressed in breast and brain cancers (gliomas). All glypicans contain an N-terminal signal peptide and a hydrophobic domain in their C-terminal region which is required for attachment of the GPI anchor. The amino acid sequences of the six vertebrate glypican family members vary from 17% to 63% identity. The location of 14 cysteine amino acids is conserved between the glypicans, suggesting the existence of a highly similar three-dimensional structure. Heparan sulfate glycosaminoglycan chains are attached at the 50 amino acids at the C-terminal end of the protein, near the anchor and the cell membrane. Glypican functions as coreceptor for a variety of growth factors. Glypican-1 has been shown to interact with SLIT2.

### Overview

Product Name	Anti-Glypican-1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Glypican-1 Antibody (Catalog # A03871). Tested in ELISA, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	P35052

### Technical Details

Immunogen	Anti-Glypican-1 protein A purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human glypican-1 protein.
Predicted Reactive Species	Chimpanzee

Isotype	IgG
Form	Liquid (sterile filtered)
Concentration	2.0 mg/mL by UV absorbance at 280 nm
Purification	Anti-Glypican-1 was protein A purified from monospecific antiserum by immunoaffinity chromatography using protein A coupled to agarose beads. This antibody is specific for human glypican-1 protein. A BLAST analysis was used to suggest partial cross-reactivity with glypican from rat, mouse, Macaque, dog, cattle, and opossum sources based on 100 - 88% homology with the immunizing sequence. Cross-reactivity with glypican from other sources has not been determined.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>ELISA: 1:10,000 - 1:50,000</p> <p>IHC: 10µg/mL</p> <p>WB: 1:500- 1:2,000</p>

## Anti-Glypican-1 Antibody (A03871) Images




Figure 1. IHC analysis of GPC1 using anti-GPC1 antibody (A03871).

GPC1 was detected in paraffin-embedded section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-GPC1 Antibody (A03871) overnight at 4°C. Biotinylated goat anti Rabbit IgG antibody was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Figure 2. Western blot analysis of GPC1 using anti-GPC1 antibody (A03871).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GPC1 antigen affinity purified polyclonal antibody (Catalog # A03871) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-Rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # SA1022) with Tanon 5200 system. A specific band was detected for GPC1.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Glypican-1 Antibody