

Anti-C/EBP Zeta (D24) CEBPZ Antibody

Catalog Number: A04074

About CEBPZ

May play a role as endocytotic lectin receptor displaying calcium-dependent lectin activity. Internalizes glycosylated ligands from the extracellular space for release in an endosomal compartment via clathrin-mediated endocytosis. May be involved in plasminogen activation system controlling the extracellular level of PLAUR/PLAU, and thus may regulate protease activity at the cell surface. May contribute to cellular uptake, remodeling and degradation of extracellular collagen matrices. May play a role during cancer progression as well as in other chronic tissue destructive diseases acting on collagen turnover. May participate in remodeling of extracellular matrix cooperating with the matrix metalloproteinases (MMPs).

Behrendt N., J. Biol. Chem. 275:1993-2002(2000). Sheikh H., J. Cell Sci. 113:1021-1032(2000). Howard M.J., J. Biol. Chem. 277:32320-32331(2002).<

Overview

Product Name	Anti-C/EBP Zeta (D24) CEBPZ Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-C/EBP Zeta (D24) CEBPZ Antibody catalog # A04074. Tested in WB,IHC applications. This antibody reacts with Human,Mouse,Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q03701

Technical Details

Immunogen	Synthesized peptide derived from human Secretogranin-1
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	lgG
Form	Liquid
Concentration	1 mg/ml





Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:500-1:1000 IHC: 1:50-1:200

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-C/EBP Zeta (D24) CEBPZ Antibody