

Anti-ADAMTS9 Antibody Picoband™

Catalog Number: A04639-1

About ADAMTS9

A disintegrin and metalloproteinase with thrombospondin motifs 9 is an enzyme that in humans is encoded by the ADAMTS9 gene. This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. Members of the ADAMTS family have been implicated in the cleavage of proteoglycans, the control of organ shape during development, and the inhibition of angiogenesis. This gene is localized to chromosome 3p14.3-p14.2, an area known to be lost in hereditary renal tumors. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar proteolytic processing.

Overview

| | |
|----------------------|---|
| Product Name | Anti-ADAMTS9 Antibody Picoband™ |
| Reactive Species | Human, Rat |
| Description | Boster Bio Anti-ADAMTS9 Antibody Picoband™ catalog # A04639-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Rat. |
| Application | ELISA, Flow Cytometry, WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ . |
| Storage Instructions | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing. |
| Host | Rabbit |
| Uniprot ID | Q9P2N4 |

Technical Details

| | |
|-------------------------------|---|
| Immunogen | E.coli-derived human ADAMTS9 recombinant protein (Position: F288-H1853). |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. |

| | |
|---------------------|---|
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.25-0.5 ug/ml, Human, Rat</p> <p>Flow Cytometry, 1-3 ug/1x10⁶ cells, Human</p> <p>Direct ELISA, 0.1-0.5 ug/ml, Human</p> |

Anti-ADAMTS9 Antibody Picoband™ (A04639-1) Images




Figure 1. Western blot analysis of ADAMTS9 using anti-ADAMTS9 antibody (A04639-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human SH-SY5Y whole cell lysates,
Lane 2: rat H9C2(2-1) whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ADAMTS9 antigen affinity purified polyclonal antibody (Catalog # A04639-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ADAMTS9 at approximately 216 kDa. The expected band size for ADAMTS9 is at 216 kDa.




Figure 2. Flow Cytometry analysis of THP-1 cells using anti-ADAMTS9 antibody (A04639-1).

Overlay histogram showing THP-1 cells stained with A04639-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ADAMTS9 Antibody (A04639-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 3. Flow Cytometry analysis of U2OS cells using anti-ADAMTS9 antibody (A04639-1).

Overlay histogram showing U2OS cells stained with A04639-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ADAMTS9 Antibody (A04639-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ADAMTS9 Antibody