

Anti-PUMA/BBC3 Antibody Picoband™

Catalog Number: A04899-3

About BBC3

The p53 upregulated modulator of apoptosis, or PUMA, is a pro-apoptotic member of the Bcl-2 protein family. The PUMA gene is located at 19q. PUMA transcript is contained within 4 exons, with the presumptive initiation codon in exon 2. The predicted 193-amino acid PUMA protein shares 91% amino acid identity with the murine sequence. Bcl-2 family members can form hetero- or homodimers, and they act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The expression of PUMA is regulated by the tumor suppressor p53, and PUMA has been shown to be involved in p53-mediated apoptosis. Additionally, PUMA encodes 2 BH3 domain-containing proteins, PUMA-alpha and PUMA-beta, that are produced through the use of an alternative first exon and are induced in cells following p53 activation. Furthermore, PUMA couples the nuclear and cytoplasmic proapoptotic functions of p53.

Overview

Product Name	Anti-PUMA/BBC3 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-PUMA/BBC3 Antibody Picoband™ catalog # A04899-3. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9BXH1

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human PUMA/BBC3, which shares 95.7% amino acid (aa) sequence identity with mouse and rat BBC3.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5 ug/ml/ml, Human

Anti-PUMA/BBC3 Antibody Picoband™ (A04899-3) Images




Figure 1. Western blot analysis of PUMA/BBC3 using anti-PUMA/BBC3 antibody (A04899-3).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human K562 whole cell lysates,

Lane 3: human Hela whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PUMA/BBC3 antigen affinity purified polyclonal antibody (Catalog # A04899-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PUMA/BBC3 at approximately 18 kDa. The expected band size for PUMA/BBC3 is at 18 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PUMA/BBC3 Antibody