

## Anti-Flt3 ligand/Flt3lg Antibody Picoband™

Catalog Number: A04987-1

### About Flt3lg

FLT3LG (FMS-Related Tyrosine Kinase 3 Ligand) also called FLT3 LIGAND, FL or FLT3L, is a protein which in humans is encoded by the FLT3LG gene. FLT3LG controls the development of DCs and is particularly important for plasmacytoid DCs and CD8-positive classical DCs and their CD103-positive tissue counterparts. Flt3 ligand (FL) is a hematopoietic four helical bundle cytokine. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors. Lyman et al. (1993) found that Flt3 ligand stimulated proliferation of hematopoietic progenitor cells isolated from mouse fetal liver or adult mouse bone marrow. Hannum et al. (1994) concluded that FL enhances the response of stem and primitive progenitor cells to other growth factors to generate all myeloid lineages except erythroid cells. Assays of C-terminally truncated FL proteins confirmed that the N-terminal half conferred FL activity. Depletion of the Pi3k -Mtor negative regulator Pten facilitated Flt3l-driven DC development in culture.

### Overview

|                      |                                                                                                                                                                                                       |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name         | Anti-Flt3 ligand/Flt3lg Antibody Picoband™                                                                                                                                                            |
| Reactive Species     | Mouse, Rat                                                                                                                                                                                            |
| Description          | Boster Bio Anti-Flt3 ligand/Flt3lg Antibody Picoband™ catalog # A04987-1. Tested in ELISA, WB applications. This antibody reacts with Mouse, Rat.                                                     |
| Application          | ELISA, WB                                                                                                                                                                                             |
| Clonality            | Polyclonal                                                                                                                                                                                            |
| Formulation          | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.                                                                                                                             |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit                                                                                                                                                                                                |
| Uniprot ID           | A0A0G2K6K8                                                                                                                                                                                            |

### Technical Details

|                               |                                                                                                 |
|-------------------------------|-------------------------------------------------------------------------------------------------|
| Immunogen                     | E. coli-derived rat Flt3 ligand recombinant protein (Position: T28-Q189).                       |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity              | No cross-reactivity with other proteins.                                                        |
| Isotype                       | Rabbit IgG                                                                                      |
| Form                          | Lyophilized                                                                                     |

|                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Concentration       | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.                                                                                                                                                                                                                                                                                                                                                                                                       |
| Purification        | Immunogen affinity purified.                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml</p> <p>Direct ELISA, 0.1-0.5ug/ml</p> |

## Anti-Flt3 ligand/Flt3lg Antibody Picoband™ (A04987-1) Images




Figure 1. Western blot analysis of Flt3 ligand using anti-Flt3 ligand antibody (A04987-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: mouse spleen tissue lysates,

Lane 2: mouse kidney tissue lysates,

Lane 3: rat skeletal muscle tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes.

Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Flt3 ligand antigen affinity purified polyclonal antibody (Catalog # A04987-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Flt3 ligand at approximately 26KD. The expected band size for Flt3 ligand is at 26KD.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Flt3 ligand/Flt3lg Antibody™