

Anti-Musashi 1/Msi1 Antibody Picoband™

Catalog Number: A05052-2

About MSI1

RNA-binding protein Musashi homolog 1 is a protein that in humans is encoded by the MSI1 gene. This gene encodes a protein containing two conserved tandem RNA recognition motifs. Similar proteins in other species function as RNA-binding proteins and play central roles in posttranscriptional gene regulation. Expression of this gene has been correlated with the grade of the malignancy and proliferative activity in gliomas and melanomas. A pseudogene for this gene is located on chromosome 11q13.

Overview

Product Name	Anti-Musashi 1/Msi1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Musashi 1/Msi1 Antibody Picoband™ catalog # A05052-2. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O43347

Technical Details

Immunogen	E.coli-derived human Musashi 1/Msi1 recombinant protein (Position: Q33-S347).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

	kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25-0.5ug/ml, Human Flow Cytometry, 1-3ug/1x10 ⁶ cells, Human Direct ELISA, 0.1-0.5ug/ml, Human
--	--



Anti-Musashi 1/Msi1 Antibody Picoband™ (A05052-2) Images




Figure 1. Western blot analysis of MSI1 using anti-MSI1 antibody (A05052-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human U2OS whole cell lysates,

Lane 2: human T-47D whole cell lysates,

Lane 3: human COLO-320 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MSI1 antigen affinity purified polyclonal antibody (Catalog # A05052-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MSI1 at approximately 39KD. The expected band size for MSI1 is at 39KD.




Figure 2. Flow Cytometry analysis of A549 cells using anti-MSI1 antibody (A05052-2).

Overlay histogram showing A549 cells stained with A05052-2 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MSI1 Antibody (A05052-2, $1ug/1x10^6$ cells) for 30 min at 20° C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5- $10ug/1x10^6$ cells) was used as secondary antibody for 30 minutes at 20° C. Isotype control antibody (Green line) was rabbit IgG ($1ug/1x10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.