

## **Anti-EIF3K Antibody**

Catalog Number: A07789

#### **About EIF3K**

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. HAMAP-Rule MF\_03010

Mayeur G.L., Eur. J. Biochem. 270:4133-4139(2003).

Nawa G., Submitted (OCT-1998) to the EMBL/GenBank/DDBJ databases.

Ye M., Submitted (JUL-1998) to the EMBL/GenBank/DDBJ databases.

#### Overview

Product Name	Anti-EIF3K Antibody
Reactive Species	Human
Description	Boster Bio Anti-EIF3K Antibody (Catalog # A07789). Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UBQ5

#### **Technical Details**

Immunogen	Synthesized peptide derived from internal of human EIF3K.
Predicted Reactive Species	Chimpanzee, Drosophila, Macaque
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitopespecific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blotting: 1:500~1:3000



### Anti-EIF3K Antibody (A07789) Images



Western blot analysis of extracts from HeLa cells, using EIF3K antibody A07789.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-EIF3K Antibody